

[Home](#) » Seresto™ Small or Large Dog (BAYER HEALTHCARE LLC)



BAYER HEALTHCARE LLC
Animal Health Division
P.O. BOX 390, SHAWNEE MISSION, KS, 66201-0390

Customer Service Tel.: 800-633-3796

Customer Service Fax: 800-344-4219

Website: www.bayer-ah.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

SERESTO™ Small or Large Dog
Bayer Animal Health

SERESTO™ small dog

For 8-month prevention and treatment of ticks and fleas on small dogs and puppies 7 weeks of age and older and up to 18 lbs. (8 kg)

SERESTO™ large dog

For 8-month prevention and treatment of ticks and fleas on large dogs and puppies 7 weeks of age and older and above 18 lbs. (8 kg)

- Repels and kills ticks for 8 months, including Deer ticks (vector of Lyme disease and anaplasmosis), American dog ticks (vector of Rocky Mountain spotted fever and ehrlichiosis), Brown dog ticks (vector of ehrlichiosis, anaplasmosis, bartonellosis, canine hemoplasmosis, babesiosis, and Rocky Mountain spotted fever), and Lone Star ticks (vector of ehrlichiosis).
- Prevents tick infestations within 48 hours after application.
- Re-infesting ticks are repelled and/or killed as quickly as 6 hours.
- Quickly kills fleas on dogs within 24 hours and continues to prevent infestations for 8 months.
- Treatment with Seresto kills fleas and may reduce the risk of tapeworm infections (*Dipylidium caninum*), bartonellosis (*Bartonella vinsonii*, *Bartonella* spp.), and rickettsiosis (*Rickettsia felis*), which are types of canine vector borne diseases).
- Re-infesting fleas are killed within 2 hours with protection against further flea infestation lasting 8 months.
- Kills fleas before they lay eggs.
- Seresto aids in the control of flea larvae in the dog's surroundings following contact with a Seresto-treated dog.
- Kills chewing lice for one month.
- Aids in the treatment and control of Sarcoptic mange.
- No need to remove collar when your pet goes swimming or gets bathed.
- Bayer Polymer Matrix ensures that both active ingredients are slowly and continuously released in low concentrations.
- Make sure to replace the collar after 8 months for year-round pest prevention.
- Light reflectors for added visibility in the evening and at night.

Active Ingredients:	% by Weight
Flumethrin*	4.5%
Imidacloprid	10.0%
OTHER INGREDIENTS	85.5%
Total	100%

* Trans Z-1/trans Z-2 ratio: max 66% trans Z-1 and min 34% trans Z-2

READ THE ENTIRE LABEL BEFORE USE

DO NOT OPEN UNTIL READY TO USE

DO NOT LET CHILDREN PLAY WITH THIS COLLAR OR REFLECTORS

KEEP OUT OF REACH OF CHILDREN

CHOKING HAZARD - CONTAINS SMALL PARTS

HOT LINE NUMBER
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
• For CONSUMER QUESTIONS, call 1-800-255-6826.
• For medical emergencies involving HUMANS, call 1-800-422-9874.
• For medical emergencies involving ANIMALS, call 1-800-422-9874.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not contaminate feed or food.

HOW TO APPLY

One collar per animal to be fastened around the neck. Remove collar from protective bag directly before use. Unroll collar and make sure that there are no remnants from the plastic connectors inside the collar.

Insert end of the collar through the buckle. Adjust the collar around the animal's neck without tightening it too tight (as a guide, it should be possible to insert 2 fingers between the collar and the neck). Pull excess collar through the loop(s) and cut off any excess length extending 1 inch (2 cm) beyond the loop(s).

After application of the collar, three reflector clips may be fixed permanently to the collar to increase the animal's visibility in the dark. Take the clips out of the bag. The clips should be evenly distributed on the non-overlapping part of the collar. The clips are correctly applied when a clicking sound is heard.

For safety reasons once the reflectors are fixed to the collar the clasp closes permanently and cannot be re-opened.



The collar should be worn continuously for the 8 month protection period. Check periodically and adjust fit if necessary, especially when puppies are rapidly growing.

This collar is designed with a safety-closure mechanism. In the unlikely event of a dog being trapped, the dog's own strength is sufficient to widen the collar to allow for quick release.

Replace the collar after 8 months for optimal tick and flea protection.

PRODUCT INFORMATION

Kills fleas which may cause flea allergy dermatitis (FAD) or flea bite hypersensitivity (FBH).

Seresto kills existing fleas on dogs within 24 hours. Reinfesting fleas are killed within 2 hours with protection against further flea infestation lasting eight (8) months. Pre-existing pupae in the environment may continue to emerge for six (6) weeks or longer depending upon the climatic conditions. Seresto aids in control of flea larvae in the dog's surroundings following contact with a Seresto-treated dog.

Ticks already on the dog prior to treatment may not be killed immediately after collar application and may remain attached and visible. At the time of application remove ticks already on the dog. The prevention of infestations with new ticks starts within 48 hours after application of the collar.

Re-infesting ticks are repelled and/or killed as quickly as 6 hours.

Kills lice on dogs for one month.

Aids in the treatment and control of Sarcoptic mange.

Seresto is water-resistant and remains effective following a shampoo treatment, swimming or after exposure to rain or sunlight. Under normal conditions, effectiveness lasts for 8 months. In order to maintain an eight-month duration, dogs must not be bathed more than once per month. For dogs that swim once a month or more, the control duration is reduced to 5 months.

Occasionally slight itching may be observed in animals that are not used to wearing collars in the first few days after fitting. Ensure that the collar is not fitted too tightly. Slight hair loss and mild skin reactions due to the mechanical irritation of the collar may occur at the application site which usually recover within 1 or 2 weeks without the need for collar removal. If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinarian.

In case of loss of collar, a new collar may be applied immediately.

Do not attach a leash to the collar as this may cause breakage.

Keep the collar in the bag and in the outer packaging until use. As with any pesticide product, do not allow small children to play with the collar or reflectors, or to put them into their mouths. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and cold water after fitting the collar. People with sensitivity reactions to the ingredients of the collar should avoid contact with the collar. Choking hazard. Contains small parts. Do not place collar or reflectors in mouth. Not intended for use on humans.

For external use on dogs only. Do not use on other animals. Do not use on puppies under seven weeks of age. Do not get this product in dog's eyes or mouth. As with any product, consult your veterinarian before using this product on debilitated, aged, breeding, pregnant, or nursing animals. Individual sensitivities, while rare, may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water. If signs persist, or become more severe, consult a veterinarian immediately. If your animal is on medication, consult your veterinarian before using this or any other product. If your dog is experiencing an adverse event, contact your veterinarian and call 1-800-422-9874.

The Bayer Polymer Matrix System ensures that both active ingredients are slowly and continuously released in low concentrations from the collar towards the animal. This avoids peak concentrations and ensures that concentrations of both active ingredients are present in the dog's haircoat during the entire efficacy period. The active ingredients spread from the site of direct contact over the entire skin surface.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original unopened container in a cool, dry place that is inaccessible to children.

Container Disposal: Dispose of container, pouch and expired collar in trash.

LIMITED WARRANTY AND LIMITATION ON DAMAGES

Bayer HealthCare LLC, Animal Health Division, warrants that this material conforms to the chemical description on the label. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, and no agent of Bayer is authorized to do so except in writing with a specific reference to this warranty. To the extent consistent with applicable law any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc. The reflector clips contained herein are only meant to be used in combination with the specific collar and have been tested only for such use and purpose. To the extent consistent with applicable law, Bayer excludes liability for any misuse and use other than in combination with the specific collar as described in the packaging insert.

EPA Reg. No. 11556-155

EPA Est. No. 11556-DEU-1

Manufactured For Bayer HealthCare LLC, Animal Health Division, P.O. Box 390, Shawnee Mission, Kansas 66201 USA

Made in Germany

81080992

81080917

17560

Regulations for product use are established by country. Information contained on this site pertains only to the United States of America, and is not intended to provide adequate information for product use. Before using or dispensing any product, read and carefully observe the label directions.

Copyright 2013 - North American Compendiums